## Track

Abstract submission: Clinical

Topic

Clinical track: Head and Neck

Presentation preference

Oral presentation

Abstract title

Deep learning auto contouring of OAR for HN radiotherapy: a blinded evaluation by clinical experts

Vincent Grégoire1, Pierre Blanchard2, Albina Allajbej1, Claire Petit2, Nicolas Milhade1, France Nguyen2, Sofia Bakkar2, Geoffroy Boulle2, Edouard Romano2, Wael Zrafi2, Aurélien Lombard3, Eugénie Ullmann3, Nikos Paragios3, Eric Deutsch2,4, Charlotte Robert4,5.

1Léon Bérard Cancer Center- University of Lyon, Department of radiotherapy, Lyon, France.

2Gustave Roussy - Paris-Saclay University, Department of radiotherapy, Villejuif, France.

3Therapanacea, Radiotherapy, Paris, France.

4U1030 Molecular Radiotherapy, Paris-Sud University - Gustave Roussy - Inserm - Paris-Saclay University, Villejuif, France.

5Gustave Roussy - Paris-Saclay University, Department of Medical Physics, Villejuif, France.

**Purpose or Objective** 

Coutouring is one of the most time-consuming steps in the radiotherapy workflow. The accuracy of high precision image-guided delivery techniques is hampered by potential deviation in target and normal tissue volume delineation. Artificial intelligence can accelerate organs-at-risk (OAR) delineation and homogenize volume definition. This study aims at evaluating a commercial solution that explores an ensemble of anatomically preserving deep-learning-based networks in two radiotherapy sites with expertise in head-and-neck cancers.

## Material and Methods

ART-Plan is a CE-marked solution for automatic annotation of OAR harnessing a unique combination of anatomically preserving and deep learning annotation concept. In average 6,600 samples were used for training per organ after data augmentation. Evaluation of the software was performed in two phases. In phase I, 100 patients were retrospectively selected in centres 1 and 2. For each patient, ART-Plan was used to generate full annotation of 15 OAR. Using a random selection, the contours generated from ART-Plan were blended with the ones corresponding to the clinical expert. For the whole cohort, 50% of the structures were the ones produced from the ART-Plan and the remaining ones from the expert in a random manner. Each contour was then scored by 5 experts, as A/clinically acceptable, B/clinically acceptable after minor corrections, C/not acceptable. The second phase of evaluation refers to the time gain between a fully manual delineation and one targeting to correct the outcomes of automatic contouring. This was done for 50 patients (25 patients from each centre) with respect to a full annotation of the 15 structures that were considered also in phase I.

## Results

96% of all manual contours were classified as clinically relevant (75% and 21% for A and B categories respectively). Values were equal to 98% for automatic contouring (56 % and 39 % for A and B respectively). Spinal cord and oral cavity obtained better scores for automatic contouring than for manual contouring (77 % and 89 % of score A for spinal cord and oral cavity versus 65 % and 64 % for manual contouring). On the contrary, optical nerves and mandibular glands were more difficulty delineated by the automatic solution. Inter-observer variability was high between experts. Average consensus for phase I was 63% between experts ranging from 53% to 77%. The time observed to correct the automated contours was significantly inferior to the time required to generate contours fully manually. In average, 2 minutes were needed to correct the contours after auto-segmentation versus 30 minutes for manual delineation.

## Conclusion

This is the first blinded, multicentric, and random back to back evaluation of an automated engine for delineation in HN tumours. The results are highly promising suggesting that this deep-learning based method should contribute to provide clinically acceptable OAR delineation. Further evaluation is ongoing to quantify the dosimetric impact of the variations observed.

Corporate sponsored research:

PhD grant

Keyword

Delineation